

Extended closed inflationary universes

Sergio del Campo*

*Instituto de Física, Pontificia Universidad Católica de Valparaíso,
Av. Brasil 2950, Casilla 4059, Valparaíso, Chile.*

Ramón Herrera†

Departamento de Ciencias Físicas, Universidad Andrés Bello, Av. República 237, Santiago, Chile.

(Dated: February 7, 2008)

In this paper we study a type of model for closed inflationary universe models using the Jordan-Brans-Dicke theory. Herein we determine and characterize the existence of $\Omega > 1$, together with the period of inflation. We have found that our model, which takes into account a Jordan-Brans-Dicke type of theory, is less restrictive than the one used in Einstein's theory of general relativity. Our results are compared to those found in Einstein's theory of Relativity.

PACS numbers: 98.80.Jk, 98.80.Bp

I. INTRODUCTION

The existence of Doppler peaks and their respective localization tend to confirm the inflationary model, which is associated with a flat universe with $\Omega = 1.02 \pm 0.02$, as corroborated by the existence of an almost scale invariant power spectrum, with $n_s = 0.97 \pm 0.03$ [1].

The recent temperature anisotropy power spectrum measured with the Wilkinson Microwave Anisotropy Probe (WMAP) at high multipoles is in agreement with an inflationary Λ -dominated CDM cosmological model. However, the low order multipoles have lower amplitudes than expected from this cosmological model[2]. Perhaps, these amplitudes may indicate the need for new physics. Speculations for explaining this discrepancy has been invoked in the sense that the low quadrupole observed in the CMB is related to the curvature scale[3].

Due to this it may be interesting to consider other inflationary universe models where spatial curvature is taken into account. In fact, it is interesting to check if the flatness, in the curvature as well as in the spectrum are indeed reliable and robust predictions of inflation[4].

In the context of an open scenario, it is assumed that the universe has a lower-than-critical matter density and, therefore, a negative spatial curvature. Several authors [5, 6, 7, 8], following previous speculative ideas [9, 10], have proposed possible models, in which open universes may be realized, and their consequences, such as density perturbations, have been explored [11]. The only available semi-realistic model of open inflation with $1 - \Omega \ll 1$ is rather unpleasant; since it requires a finely-tuned potential of very peculiar shape [8, 12]. Very recently the possibility to create an open universe from the perspective of the brane-world scenarios has been considered [13].

The possibility of having inflationary universe models

with $\Omega > 1$ has also been considered[4, 14, 15]. Particularly, this case has been marginally indicated by the WMAP recent observations[3].

In this paper we study inflationary closed universe models in which a Jordan-Brans-Dicke (JBD) theory is taken into account[16]. Following the line described by Linde[4], we considered two situations: in the first, the inflaton potential $V(\sigma)$ is constant. More precisely, in this case the model is based on a step-like effective potential in which $V(\sigma) = 0$ at $\sigma < 0$ and $V(\sigma) = V = \text{const}$ for the range of σ given by $0 < \sigma < \sigma_o$, where σ_o is its initial value. Close to this value ($\sigma = \sigma_o$) we will assume that the potential sharply rises to infinitely large values. In a universe with $\Omega > 1$, and in order to make inflation short, one could consider that the effective potentials at the Planck density are extremely steep. In general, such a universe will not typically enter the inflationary regime and will collapse quickly. This suggests that most of the universes described by the models that use extremely steep potentials will not inflate at all, or will expand by a factor much less than 60 e-folds. In this way, most of the universe will be short living. We like to address this particular problem here; the other effective potential that we are considering is a chaotic potential of the form $V(\sigma) = \lambda_n \sigma^n / n$. If we agree with the probability of creation of an inflationary universe, it is suggested that it is much more natural for the universe to be created with a density very closed to the Planck density.

In the case of Einstein's theory of general relativity with a constant scalar inflaton potential $V(\sigma)$, the inflaton field σ stops before it reaches the value $\sigma = 0$, whose value is necessary for ending inflation, and hence inflation would continue forever[4]. Differently, in a JBD type of theory, it is possible to solve this (graceful exit) problem. As we will see, the JBD parameter ω and the initial condition of the JBD field make it possible that the inflaton field reaches the value $\sigma = 0$ and thus solving the graceful exit problem.

The structure of the paper is as follows: In Sec. II we present the action and equations for the Jordan-Brans-Dicke theory. In Sec. III we determine the characteristics

*E-mail address: sdelcamp@ucv.cl

†E-mail address: rherrera@unab.cl

of the closed inflationary universe models that are produced with a potential $V(\sigma) = \text{constant}$. In Sec. IV we determine closed inflationary universe models for chaotic potentials of the form $V(\sigma) = \lambda_n \sigma^n / n$. We give an estimation of the scalar density perturbation $\delta\rho/\rho$. At a different level, our results are compared with those obtained with Einstein's theory of gravity. Finally, conclusions and some discussions are presented in Sec. V.

II. THE COSMOLOGICAL EQUATIONS IN JBD THEORY

We consider the effective action S given by

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} \varepsilon \phi^2 R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \mathcal{L}(\sigma) \right], \quad (1)$$

where

$$\mathcal{L}(\sigma) = \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma - V(\sigma),$$

and R is the Ricci scalar curvature, ϕ is the JBD scalar field, and ε is a dimensionless coupling constant that, in terms of the JBD parameter $\omega = \frac{1}{4\varepsilon}$. $V(\sigma)$ is an effective scalar potential associated with the inflaton field, σ .

The Friedmann-Robertson-Walker metric is described by

$$ds^2 = dt^2 - a(t)^2 d\Omega_k^2, \quad (2)$$

where $a(t)$ is the scale factor, t represents the cosmic time and $d\Omega_k^2$ is the spatial line element corresponding to the hypersurfaces of homogeneity, which could represent a three-sphere, a three-plane or a three-hyperboloid, with values $k=1, 0, -1$, respectively. From now on we will restrict ourselves to the case $k=1$ only.

Using the metric (2), and $k=1$, in the action (1), we obtain the following field equations:

$$\ddot{\sigma} = -3 \frac{\dot{a}}{a} \dot{\sigma} - \frac{dV}{d\sigma}, \quad (3)$$

$$\ddot{\phi} = -3 \frac{\dot{a}}{a} \dot{\phi} - \frac{\phi^2}{\phi} - \frac{1}{1+6\varepsilon} \left[\frac{\dot{\sigma}^2}{\phi} - \frac{4}{\phi} V(\sigma) \right], \quad (4)$$

$$\frac{\dot{a}^2}{a^2} + 2 \frac{\dot{a}\dot{\phi}}{a\phi} = -\frac{1}{a^2} + \frac{1}{3\varepsilon\phi^2} \left[\frac{\dot{\phi}^2}{2} + \frac{\dot{\sigma}^2}{2} + V(\sigma) \right], \quad (5)$$

and

$$\ddot{a} = 2 \frac{\dot{a}\dot{\phi}}{\phi} - \frac{a}{3\varepsilon\phi^2} \left[\dot{\phi}^2 + \frac{1+3\varepsilon}{1+6\varepsilon} \dot{\sigma}^2 - \frac{1-6\varepsilon}{1+6\varepsilon} V(\sigma) \right], \quad (6)$$

where the dots over ϕ and a denotes derivatives with respect to the time t . For convenience we will use from now on units where $c = \hbar = M_p = G^{-1/2} = 1$. Note that this set of equation reduces itself to the set of Einstein's field Eqs., in the limits $\varepsilon \rightarrow 0$ and $\varepsilon\phi^2 \rightarrow M_p^2/8\pi = 1/8\pi$.

III. CLOSED INFLATIONARY UNIVERSE MODEL WITH $V = Cte$.

Following Linde[4], let us consider for simplicity a toy model with the following step-like effective potential: $V(\sigma) = 0$ at $\sigma < 0$; $V(\sigma) = V = \text{const}$ at $0 < \sigma < \sigma_o$ (here σ_o is the initial value of the inflaton field). We shall also assume that the effective potential sharply rises to infinitely large value in a small vicinity of $\sigma = \sigma_o$. As Linde mentions in Ref.[4] this model is very well suited for the description of a certain version of an F-term hybrid inflation in supergravity. In such models the effective potential is nearly constant at small σ and it exponentially rises at large σ . We should mention that the physical background of this model is inspired by Linde's Hybrid inflation model[17]. Therein, the inflationary plateau is due to the vacuum density associated with some other so-called "waterfall" field which is coupled to the inflaton field. Here, the plateau is slightly tilted due to radiative corrections so the field will continue to roll down, and in addition the end of inflation occurs at some critical value $\sigma_c > 0$, differing from our case in which inflation ends at $\sigma = 0$. But, fundamentally, the models are rather similar. In fact, for large values of σ , supergravity corrections introduce a quartic term which grows rapidly and resembles very much the abrupt cliff mentioned above.

Suppose now that a close universe described by these models appeared "from nothing" in a state with the field $\sigma \geq \sigma_o$ at the point with $\dot{a} = 0$, $\dot{\sigma} = 0$, $\dot{\phi} = \sqrt{\beta_o} \tilde{\phi}_o$ and the potential energy density equal to V_o (with $V_o \geq V$). Here, the time derivative of the JBD scalar $\dot{\phi}$ has an initial condition that depends on β_o and $\tilde{\phi}_o$. The value of β_o permits us to establish when inflation may occur, by means of the probability of creation of an inflationary universe model. This restricts the initial value of the inflaton field, σ_o , and also, the value of $\tilde{\phi}_o$ which represents a fraction of the initial value of ϕ . We choose this particular initial value for $\dot{\phi}$ in order that, in each one of the successive intervals of the inflaton field, $\Delta\sigma$, and in each one of the successive intervals of the inflationary process, these quantities do not depend on β_o .

It will be convenient to represent V_o as

$$V_o = V^* - \Delta V^*,$$

where $V^* = Cte. = \frac{3V}{2(1+3\varepsilon)}$. Note that this quantity represents an upper bound on V_o which occurs when $\Delta V^* \rightarrow 0$. The definition of V^* in term of V and ε allows us to study the energy density in the vicinity of $\sigma = \sigma_o$ and the one established from energy conservation¹ and the

¹ In the case of GR theory:

The energy conservation $V_o + 0 = \frac{\dot{a}^2}{2} + V \Rightarrow \dot{\sigma}^2 = 2(V_o - V)$ in Eq. for $\ddot{a}$ i.e.

$$\ddot{a} = \frac{8\pi}{3} a(V - \dot{\sigma}^2) = \frac{8\pi}{3} a(3V - 2V_o)$$

equation(6). Note that this value of V^* yields the appropriate Einstein's General Relativity limit for $\varepsilon \rightarrow 0$. Here, ΔV^* is a small quantity that becomes a function of the quantities ε , V and β_o .

Then, the field σ instantly would fall down to the plateau ($V(\sigma) = V$), and its potential energy density would become converted into the kinetic energy, $\frac{\dot{\sigma}^2}{2} = \frac{(1-6\varepsilon)V}{2(1+3\varepsilon)} - \Delta V^*$, and thus the velocity of the field σ would be given with $\dot{\sigma} = -\sqrt{(1-6\varepsilon)V/(1+3\varepsilon) - 2\Delta V^*}$. Since this happens nearly instantly, one still has $\dot{a} = 0$, $\dot{\phi} = \sqrt{\beta_o \dot{\phi}_o}$ and $\dot{\sigma} = -\sqrt{(1-6\varepsilon)V/(1+3\varepsilon) - 2\Delta V^*}$ at these times. These values could be considered initial conditions when solving the set of Eqs.(3)-(6), for $V(\sigma)=V=\text{const}$ in the interval $0 < \sigma < \sigma_o$. Since the field σ instantly falls down to the plateau ($V=\text{const}$), its potential energy density is converted into kinetic energy. This causes $\dot{\sigma}$ to change from a zero value to a negative constant value. During this change, we see from the Friedmann equation that $\dot{a} = 0$ and $\phi = \text{const.} = \phi_o$. After this period, the inflaton field σ instantly falls down to the plateau and the evolution of the JBD field ϕ monotonically increases to some constant value, which we denote by ϕ_N . We fix this value to the actual value of the Planck mass times $\sqrt{\omega/2\pi}$ (recall that $\phi_N^2 = M_p^2/8\pi\varepsilon$ which becomes $\phi_N^2 = \omega/2\pi = \text{const.}$ in unit with $M_p = 1$). In Fig. 1 we show the temporal evolution of the JBD scalar field.

In order to clarify the conditions under which a closed universe could remain static, or could collapse, or even could enter an inflationary regime, it is necessary to rewrite Eq.(6) in terms of ΔV^*

$$\ddot{a} = a\eta \left[2H - \frac{1}{3\varepsilon}\eta \right] + \frac{2a(1+3\varepsilon)}{3\varepsilon\phi^2(1+6\varepsilon)}\Delta V^*,$$

where $H = \dot{a}/a$ defines the Hubble expansion rate and $\eta = \dot{\phi}/\phi = |\frac{\dot{G}}{2G}| \ll 1$ where $G = 1/8\pi\varepsilon\phi^2$ is the gravitational "constant". The present value of η is of the order of $\eta_o \leq 10^{-10}/\text{yr}$ which in Planck mass unit becomes $\sim 10^{-45}M_p$. In general, during inflation we have $H \sim \sqrt{V}/M_p$ with $\sqrt{V} \sim 10^{-5}M_p$ and we assume that during inflation $H > \eta/6\varepsilon$. This assumption is justified, since during inflation $H_{inf} \sim 10^{-5}M_p$. In our models, we take $\eta_{inf} \sim 10^{-11}M_p$, when we resolve the field equations for the different values of the JBD parameter ω ,




FIG. 1: For the model $V = \text{const}$ we have plotted the JBD field $\phi(t)$ as a function of the time t , for two different values of the JBD parameter, $\omega = 500$ and $\omega = 2000$. We have assumed the constant V to be equal to one.

initial values of the JBD field and the time derivatives of the JBD scalar $\dot{\phi}$. This approximation coincides with the approach used by the author in Ref.[18], when he studies inflation in a JBD theory. In this form, the above equation takes the form $\ddot{a} \simeq 2a\eta H + \frac{2a(1+3\varepsilon)}{3\varepsilon\phi^2(1+6\varepsilon)}\Delta V^*$.

In the particular case of $\Delta V^* = 0$, different possibilities exist as opposed to what happens in Einstein's GR theory. In the case of $\eta H > 0$, $\ddot{a} > 0$ and $3\dot{a}/a > 0$, the value of $\dot{\sigma}$ decreases, and the universe enters an inflationary phase. In the case of $\eta H < 0$, $\ddot{a} < 0$ and $3\dot{a}/a < 0$, the universe collapses. If $H\eta + \frac{(1+3\varepsilon)}{3\varepsilon\phi^2(1+6\varepsilon)}\Delta V^* = 0$, the acceleration of the scale factor $\ddot{a} = 0$ and the universe remains static, as occurs in Einstein's GR theory.

For $H\eta + \frac{(1+3\varepsilon)}{3\varepsilon\phi^2(1+6\varepsilon)}\Delta V^* < 0$, the term $3\dot{a}/a < 0$, which makes the motion of σ even faster with the consequence of a rapidly collapsing universe.

In the case of $H\eta + \frac{(1+3\varepsilon)}{3\varepsilon\phi^2(1+6\varepsilon)}\Delta V^* > 0$, which means $\ddot{a} > 0$, we get $3\dot{a}/a > 0$. The field σ rapidly decreases, and thus an inflationary phase is generated.

Note also that, in the regime where $V=\text{const}$, the scalar field σ satisfies equation

$$\ddot{\sigma} = -3\frac{\dot{a}}{a}\dot{\sigma}, \quad (7)$$

Static universe $\ddot{a} = 0 \Rightarrow V_o = 3V/2$; Collapses $\ddot{a} < 0 \Rightarrow V_o > 3V/2$; Inflation $\ddot{a} > 0 \Rightarrow V_o < 3V/2$. In general $V_o = 3V/2 - \Delta V$ with $\Delta V = 0, < 0, > 0$.

In the case of JBD Theory (analogy):

The energy conservation $V_o + 0 = \frac{\dot{\sigma}^2}{2} + V \Rightarrow \dot{\sigma}^2 = 2(V_o - V)$ in Eq.(6), we find

$$V_o = \frac{3V}{2(1+3\varepsilon)} - \Delta V^* = V^* - \Delta V^*.$$

which implies that

$$\dot{\sigma}(t) = \dot{\sigma}_o \left(\frac{a_o}{a(t)} \right)^3. \quad (8)$$

Here $\dot{\sigma}_o$ is the initial velocity of the field σ , immediately after it rolls down to the flat part of the potential. But once expansion of the universe begins, this expansion follows a simple power-law (as opposed to the GR case where $a \sim e^{Ht}$, with $H = \sqrt{8\pi V/3}$, see Ref.[4]). The effect of the JBD scalar field in this model is reflected in the change of the slope of the scalar field σ , even if its form remains the same, as shown in Figs. 2 and 3. We could describe a power-law regime, (which occurs if we neglect $\dot{\phi}/\phi$ as compared with H , $\dot{\phi}$ as compared with $3H\dot{\phi}$ and $\dot{\phi}^2$, $\dot{\sigma}^2$ as compared with the effective potential $V(\sigma)$, and the term $1/a^2$ is rapidly diluted) and possible solutions would be given by [18]

$$\frac{a(t)}{a_o} = \left[\frac{\phi(t)}{\phi_o} \right]^\gamma = (\alpha t + 1)^\gamma, \quad (9)$$

where

$$\gamma = \frac{(1+6\varepsilon)}{4\varepsilon}; \quad \alpha = \frac{4}{(1+6\varepsilon)\phi_o^2} \sqrt{\frac{\varepsilon\phi_o^2 V}{3}},$$

in which ϕ_o is the initial value of the JBD.

The resulting equation for the scale factor reads as follows:

$$\ddot{a} = 2\frac{\dot{a}\dot{\phi}}{\phi} - \frac{a\dot{\phi}^2}{3\varepsilon\phi^2} + \frac{2aV\beta(t)}{3\varepsilon\phi^2}. \quad (10)$$

At this point, we will introduce a small time-dependent function defined by

$$\beta(t) = \frac{1}{2V(1+6\varepsilon)}[(1-6\varepsilon)V - (1+3\varepsilon)\dot{\sigma}^2] \ll 1. \quad (11)$$

Here, we would like to make a simple analysis of the solutions to Eq.(10) for $\beta(0) \equiv \beta_o \ll 1$, in which

$$\begin{aligned} \beta_o &= \frac{1}{2V(1+6\varepsilon)}[(1-6\varepsilon)V - (1+3\varepsilon)\dot{\sigma}_o^2] = \\ &= \frac{(1+3\varepsilon)\Delta V^*}{(1+6\varepsilon)V} \ll 1, \end{aligned} \quad (12)$$

when $a(t)$ increases enough and an inflationary regime settles in, the inflaton scalar field σ gradually stops moving. From Eq. (8) together with Eq. (9) we write $\dot{\sigma} = \dot{\sigma}_o(\alpha t + 1)^{-3\gamma}$, and σ moves a finite distance, which in our case becomes

$$\Delta\sigma_{\text{inf}} = \frac{\dot{\sigma}_o}{\alpha(3\gamma-1)} \approx -\frac{(1+6\varepsilon)}{(3+14\varepsilon)} \sqrt{\frac{(1-6\varepsilon)3\varepsilon\phi_o^2}{(1+3\varepsilon)}}. \quad (13)$$

When $\varepsilon \rightarrow 0$ and $\varepsilon\phi_o^2 = Cte. = 1/8\pi$, we obtain $\Delta\sigma_{\text{inf}} \approx -1/(2\sqrt{6\pi})$, which coincides with the result obtained in Einstein's theory of Relativity [4].

At the beginning of the process, when $a \approx a_o$, $\phi \approx \phi_o$ and $\beta(t) \approx \beta_o$, Eq. (10) becomes

$$\ddot{a} = 2\frac{\dot{a}\dot{\phi}}{\phi_o} - \frac{a_o\dot{\phi}^2}{3\varepsilon\phi_o^2} + \frac{2a_oV\beta_o}{3\varepsilon\phi_o^2}, \quad (14)$$

and for small t the solution of Eq. (14), i.e. the scale factor $a(t)$, is found to be given by

$$a(t) \simeq a_o \left(1 + \frac{\beta_o}{3\varepsilon\phi_o^2} \left[V - \frac{\dot{\phi}_o^2}{2} \right] t^2 + O(t^3) \right). \quad (15)$$

From Eq. (8) we find that at the time interval Δt_1 given by

$$\Delta t_1 \approx \sqrt{\frac{2(1+6\varepsilon)\varepsilon\phi_o^2}{(1-6\varepsilon)(2V - \dot{\phi}_o^2)}}, \quad (16)$$

the parameter β becomes twice β_o , when consequently the inflaton field σ decreased by the amount

$$\Delta\sigma_1 \sim \dot{\sigma}(0)\Delta t_1 \approx -\sqrt{\frac{2V(1+6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)(2V - \dot{\phi}_o^2)}}. \quad (17)$$

This process continues, after a time $\Delta t_2 \approx \Delta t_1$, where now the field σ decreases by the amount $\Delta\sigma_2 \approx \Delta\sigma_1$, implying that the rate of growth of the scale factor, $a(t)$ doubles. This process finishes when $\beta(t) = (1-6\varepsilon)/2(1+6\varepsilon)$. Here, the beginning of inflation is determined by the initial value of the inflaton field given by

$$\begin{aligned} \sigma_{\text{inf}} &\approx \sigma_o + \sqrt{\frac{3(1-6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)}} \frac{(1+6\varepsilon)}{(3+14\varepsilon)} + \\ &+ \sqrt{\frac{2V(1+6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)(2V - \dot{\phi}_o^2)}} \ln \left[\frac{(1+6\varepsilon)\beta_o}{(1-6\varepsilon)} \right]. \end{aligned} \quad (18)$$

Note that this expression indicates that our results are very sensitive to the choice of particular values of ω , and ϕ_o , apart from the other parameters that enter the theory. Note that in the limit in which $\varepsilon\phi_o^2 \rightarrow 1/8\pi$ and $\varepsilon \rightarrow 0$, the above expression reduces itself to $\sigma_{\text{inf}} \approx \sigma_o + 0.1 + 0.15 \ln \beta_o$, where now, in this limit, β_o becomes $\beta_o \rightarrow (1 - \dot{\sigma}_o^2/V)/2$. Since inflation occurs in the interval $\sigma_{\text{inf}} > 0$ and $\sigma = 0$, we should obtain for the initial value of the inflaton field

$$\begin{aligned} \sigma_o &> -\frac{(1+6\varepsilon)}{(3+14\varepsilon)} \sqrt{\frac{3(1-6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)}} - \\ &- \sqrt{\frac{2V(1+6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)(2V - \dot{\phi}_o^2)}} \ln \left[\frac{(1+6\varepsilon)\beta_o}{(1-6\varepsilon)} \right]. \end{aligned} \quad (19)$$

We continue describing a model of quantum creation for a closed inflationary universe model. According to Ref.[19], the probability of the creation of a closed universe filled with a scalar field σ , with an effective potential $V(\sigma)$ in the theory of JBD for the case $\varepsilon\phi^2 \gg V^{1/2}$ is given by

$$P \sim e^{-2|S|} = \exp\left(\frac{-3(8\pi\varepsilon\phi^2)^2}{8V(\sigma)}\right). \quad (20)$$

We first estimate the conditional probability that the universe is created with an energy density equal to $V^* - (1+6\varepsilon)\beta_o V/(1+3\varepsilon)$, under the condition that its energy density V was smaller than V^* ; for the initial value of the JBD field $\phi = \phi_o$ we find

$$P \sim \exp\left(\frac{-3(8\pi\varepsilon\phi_o^2)^2}{8}\left[\frac{1}{V^* - \frac{(1+6\varepsilon)\beta_o V}{(1+3\varepsilon)}} - \frac{1}{V^*}\right]\right)$$

$$P \sim \exp\left(-\frac{(8\pi\varepsilon\phi_o^2)^2(1+6\varepsilon)^2\beta_o}{6(1+3\varepsilon)V}\right), \quad (21)$$

which implies that the process of quantum creation of an inflationary universe model is not exponentially suppressed by $\beta_o < \frac{6(1+3\varepsilon)V}{(1+6\varepsilon)^2(8\pi\varepsilon\phi_o^2)^2}$, which in turn means that the initial value of the inflaton field σ must be bounded from below

$$\sigma_o > -\sqrt{\frac{3(1-6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)}} \frac{(1+6\varepsilon)}{(3+14\varepsilon)} -$$

$$\sqrt{\frac{2V(1+6\varepsilon)\varepsilon\phi_o^2}{(1+3\varepsilon)(2V-\phi_o^2)}} \ln\left[\frac{6(1+3\varepsilon)V}{(1-36\varepsilon^2)(8\pi\varepsilon\phi_o^2)^2}\right]. \quad (22)$$

Let us consider two different cases for the values $\varepsilon\phi_o^2$ and ω .

First, suppose $V \sim 10^{-11}$, $8\pi\varepsilon\phi_o^2 = 1/2$, $\phi_o^2 = 10^{-12}$ and $\omega = 500$. In this case the probability of creation of an inflationary universe is not suppressed by $\beta_o < 2.4 \cdot 10^{-10}$. The value β_o allows us to fix the initial value of the inflaton field. In this case we have found that inflation begins when the field σ_o takes the value

$$\sigma_o > -0.08 - 0.14 \ln(2 \cdot 10^{-10}) \sim 3.0 M_p. \quad (23)$$

After the initial inflation, the field σ stops moving when it passes the distance $|\Delta\sigma_{\text{inf}}| \simeq 0.08$. However, this result is a particular value that depends on the value we assign to the parameter ω and the initial values of ϕ_o and $\dot{\phi}_o$ for a given value of the effective potential V .

Now, suppose that $V \sim 10^{-11}$, $8\pi\varepsilon\phi_o^2 = 1/8$, $\phi_o^2 = 10^{-12}$ and $\omega = 1000$. In this case, the probability of

creation of an inflationary universe is not suppressed by $\beta_o < 3.8 \cdot 10^{-9}$. And hence we found that inflation begins when the initial value of the inflaton field σ takes the value

$$\sigma_o > -0.08 - 0.15 \ln(4 \cdot 10^{-9}) \sim 2.8 M_p. \quad (24)$$

Note that in this case, the field σ moves a finite distance which becomes $|\Delta\sigma_{\text{inf}}| \simeq 0.08$. If this field stops before it reaches $\sigma=0$, then the universe inflates eternally. The same problem is found in Einstein's General Relativity model ($\Delta\sigma_{\text{inf}} \simeq -1/2\sqrt{6\pi} = \text{const.}$). However, in the context of the JBD theory, the value of $\Delta\sigma_{\text{inf}}$ is quite sensitive to the values we assign to ω , ϕ_o and $\dot{\phi}_o$. Therefore, as we will see, the problem that the universe inflated for ever disappears and the inflaton field reaches the value $\sigma=0$, for $\omega < 2000$ and for some appropriate initial conditions of the JBD field ϕ_o and $\dot{\phi}_o$. It has been claimed that the values of ω should be greater than 2000, in order that the JBD theory be consistent with the astronomical observations. In the more general scalar-tensor theory, ω can vary, depending on the scalar field ϕ . In cosmological models, based on such theories, it has been pointed out that there is generally an attractor mechanism that drives ω to ∞ in the late cosmological epochs[20]. But, at an early time in the evolution of the universe the value of $\omega(\phi)$ can be significantly different. Here, information on the different cosmological epochs may constrain the value of ω to the range $10 < \omega < 10^7$ (see Ref.[21]). Therefore, in the early universe it is possible to consider ω which is smaller than the one used at the present epoch[22].

The numerical solutions to the inflaton field $\sigma(t)$ are shown in Fig.2, for different values of the ω parameter. Note that the interval of σ_o at σ_{inf} increases when the parameter ω decreases, but its shapes remain practically unchanged. We should note here that, as long as we decrease the value of the ω parameter, the quantity $\sigma_o - \sigma_{\text{inf}}$ increases, and thus it permits $|\Delta\sigma_{\text{inf}}|$ to reach $\sigma=0$, even when they themselves are not oscillations of the inflaton field in neither of the two theories presented. If the field σ starts its motion with sufficiently small velocity, inflation begins immediately, in analogy with Einstein's GR theory. If it starts with large initial velocity $\dot{\sigma}_o$, because it falls from the value of $V(\sigma) = V^* - \Delta V^*$, the universe does not present the inflationary period at any stage.

IV. CHAOTIC INFLATION WITH $V = \lambda_n \sigma^n/n$

Let us now consider chaotic inflation universe models as a next step. In this model the main idea was to study all possible initial conditions in the universe, including those where the scalar field outside of the minimum of the scalar field potential[23]. In this sense, in order to study the evolution of the universe filled with a scalar field σ , we must somehow set the initial values of the field and its derivatives at different point in the space. Also it is needed to specify the topology of the space and its metric




FIG. 2: For the model $V = \text{const}$ and the same value of σ_o , we have plotted the inflaton field $\sigma(t)$ as a function of the time, for three different values of the JBD parameter, $\omega = 150$, $\omega = 500$ and $\omega = 2000$, respectively. Here we have assumed the initial value for the JBD field to be equal to $8\pi\epsilon\phi_o^2 = 1/2$. GR represents the same plot, however using Einstein's theory of Relativity. We have assumed the constant V to be equal one.

in a manner consistent with the initial values. In this way, this kind of model is concerned with the evolution of a universe filled with a chaotically distributed scalar field σ .

In the context of chaotic inflation, another possible realization is an axionic (shift)symmetry, whenever the inflaton is a pseudo-scalar axion. In this scheme, the inflaton potential arises due to the breaking of a (global) axionic symmetry, and it is therefore controlled by it: for instance, the coupling of the inflaton to matter does not affect the inflaton potential if the axionic symmetry is respected. This mechanism, known as natural inflation, was originally proposed in [24], and several possible implementations have been discussed in [25]. Shift symmetries also arise within string theory, and their application to inflation has been considered for instance in [25, 26].

Let us now consider an effective potential given by $V = \lambda_n \sigma^n/n$ for $\sigma < \sigma_o$, which becomes extremely steep at $\sigma > \sigma_o$. If the universe is created at σ just above σ_o , at the point $V_o > V(\sigma_o) = \lambda_n \sigma_o^n/n$, the field immediately falls down to σ_o and acquires a velocity given by $\dot{\sigma}_o^2/2 = V_o - V(\sigma_o)$.

We will denote $V^*(\sigma_o)$ as the critical initial value of

V , such that inflation still exists for the field σ falling to the point σ_o from the height $V_o < V^*$, and disappears at $V_o > V^*$. We also introduce the parameter β_o , similar to the one of the previous section. During inflation, the scale factor is given in the present situation by

$$\frac{a}{a_o} = \left(\frac{\phi}{\phi_o} \right)^\gamma = \left(1 + \frac{(\sigma_o^2 - \sigma(t)^2)}{\gamma n \epsilon \phi_o^2} \right)^{\frac{\gamma}{2}}, \quad (25)$$

and the universe expands $\exp(\gamma/2 \ln(1 + \sigma_o^2/\gamma n \epsilon \phi_o^2))$ times, when $\sigma(t) \rightarrow 0$.

We considered a model with $n = 2$, where the effective potential blows up at $\sigma > \sigma_o = 0(10)$, $\omega = 500$, $8\pi\epsilon\phi_o^2 = 1/2$, $\phi_o = 10^{-12}$ and $V = 10^{-11}$. In this case we have found that inflation begins when the σ rolls down to the point

$$\sigma_{\text{inf}} \approx \sigma_o + 0.03 + 0.14 \ln \left[\frac{(1 + 6\epsilon)\beta_o}{(1 - 6\epsilon)} \right]. \quad (26)$$

In order to achieve the 60 e-folds we first need to give the values of ϵ and ϕ_o so that we could find an initial value for the scalar field σ . For instance, if we chose $\omega=500$ and $8\pi\epsilon\phi_o^2 = 1/2$, we find that $\sigma_o = 2.3$. With these values, and $V \sim 10^{-11}$, the probability of creation of the universe is not suppressed for $\beta_o \lesssim 2.4 \cdot 10^{-10}$. Therefore one can argue that it is most probable to have $\beta_o \sim 2.4 \cdot 10^{-10}$. This means that inflation typically starts in this model at

$$\sigma_{\text{inf}} \approx \sigma_o - 3.1, \quad (27)$$

which means that inflation starts with a larger value of the scalar field σ than the one obtained in the JBD theory.

Now we consider a model with $n = 4$, where the effective potential $V = \lambda_4 \sigma^4/4$ blows up at $\sigma > \sigma_o = 0(10)$, $\omega = 500$, $8\pi\epsilon\phi_o^2 = 1/4$, $\phi_o = 10^{-12}$ and $V = 10^{-11}$. In this case we have found that inflation begins when the σ rolls down to the point

$$\sigma_{\text{inf}} \approx \sigma_o + 0.07 + 0.14 \ln \left[\frac{(1 + 6\epsilon)\beta_o}{(1 - 6\epsilon)} \right]. \quad (28)$$

As an example, we considered $\omega=500$ and $8\pi\epsilon\phi_o^2 = 1/4$, such that $\sigma_o = 3.3$ is required to obtain an universe that inflates e^{60} times. Also we have taken the values $V \sim 10^{-11}$, $\phi_o \sim 10^{-12}$ and $\beta_o \sim 9.6 \cdot 10^{-10}$. This means that the inflation field typically starts at

$$\sigma_{\text{inf}} \approx \sigma_o - 2.8. \quad (29)$$

For $\sigma_o = 10$, inflation starts at $\sigma_{\text{inf}} \sim 7.2$, the universe inflates e^{321} times and becomes flat. The universe inflates e^{60} times for $\sigma_{\text{inf}} = 2.3$, i.e. for $\sigma_o = 5.1$ and this leads to $\Omega = 1.1$. The numerical solution $\sigma(t)$ is shown in Fig.3 for some different values of the ω parameter and the same velocity, σ_o . In the same situation, we have studied

the evolution of the JBD field ϕ . We have found that this field monotonically increases towards some constant value, which is closer to the one determined by the actual value of the Planck mass (recall that $\varepsilon\phi_N^2 = 1/8\pi$), just when the inflaton scalar field σ begins to oscillate near the minimum of the effective potential, located at $\sigma \approx 0$.




FIG. 3: This plot shows the inflaton field $\sigma(t)$ as a function of the cosmological times t , for the effective potential $V(\sigma) = \lambda_4 \sigma^4/4$, either in Einstein's or in JBD theories ($\omega = 500$ and $\omega = 2000$) for the same value of σ_o . In these graphs we have assumed the constant λ_4 to be 1.

It is interesting to study our model density perturbations but, as the JBD theory affects the density perturbations, we calculated the density perturbations in our model according to Ref. [27] and plotted the function

$$\frac{\delta\rho}{\rho} \approx Cte H^2 \left[(\varepsilon\phi^2 8\pi)^{\frac{3}{2}} \frac{1}{|\dot{\sigma}|} + \frac{(1 - \varepsilon\phi^2 8\pi)\gamma\phi}{2|\dot{\phi}|} \right], \quad (30)$$

where

$$\gamma = \frac{1}{\sqrt{\omega + 3/2}},$$

and $Cte = 3/5\pi$, which corresponds to density perturbations in a flat universe, and note that equation (30) coincides with the GR, if we assume that $\varepsilon\phi^2 = 1/8\pi$. However, these density perturbations should be supplemented by several different contributions in a closed inflationary universe, which may alter the final result for

$\delta\rho/\rho$ at small N . However we will postpone a complete investigation of this problem for the time being.

Fig.(4) shows the magnitude of the density perturbations as a function of the N e-folds parameter for the model λ_4 , for $\sigma_o = 10$ and for various values of ω . Similar to the corresponding model in the Einstein's theory of GR, $\delta\rho/\rho$ has a maximum at small $N \simeq 0(7)$. Numerically, we could show that it presents a small displacement for $\omega=500$ with the maximum at $N = 0(8)$ which corresponds to a scale $\sim 10^{25}$ cm, which is similar to that obtained in the Einstein's GR theory.




FIG. 4: Scalar density perturbation for the model $V(\sigma) = \lambda_4 \sigma^4/4$ as a function of the N e-folds parameter for two different values of the JBD ω parameter. The values that we have assumed are $\omega = 500$ and $\omega = 2000$ for $\lambda_4 = 10^{-14}$ and $\sigma_o = 10$. These plots are compared with that obtained by using Einstein's GR theory, where $\delta\rho/\rho \approx Cte.H^2/|\dot{\sigma}|$.

V. CONCLUSION

In this article we study a closed inflationary universe model in a JBD theory. In Einstein's GR theory this model was studied by Linde[4]. Here, when V is constant, in the interval from σ_o to $\sigma = 0$ we found some problems. First of all, and due to the form of the potential, it is found that the universe typically either collapses very soon, or inflates forever (graceful exit problem). Secondly, after the end of inflation, which occurs when σ reaches the value $\sigma = 0$, it seems that the inflaton field does not appear to engage into oscillations. This is only

because the model is oversimplified. Actually, we should mention here that the inspiration for this model comes from the Linde's Hybrid inflationary model where the end of inflation sends the system towards a direction perpendicular to the inflaton direction in field space (the direction of the waterfall). Then, both fields begin oscillating around the true minimum of the scalar potential. Thus, the problem mentioned above does not correspond to the physical model, which is the basis for our toy-model.

In a JBD theory we have found that it is possible to successfully solve the graceful exit problem for an effective potential $V = \text{constant}$ with an extremely steep at the Planck scale. The dependence on the values of ω and the initial conditions for ϕ_o and $\dot{\phi}_o$ permit us to reach the value $\sigma = 0$, needed for solving the graceful exit problem. These conclusions are in good qualitative agreement with the result of a numerical investigation.

We have found that for some initial velocity values $\dot{\sigma}_o$, it is probable that in Einstein's GR theory the universe will have a large value of Ω ; however, in a JBD theory and for ω parameter $\omega < 2000$, it is possible to obtain $\Omega \sim 1.1$.

Also, in the case in which the potential is very sharp at large σ we have found extra ingredients in the JBD theory, when compared with its analog in Einstein's GR theory. Specifically, we obtain a short stage of inflation for a closed universe. This could be possible due to that the speed of the rolling field was very large but it could be controlled with the value of the JBD parameter ω . Similarly, the long-wavelength perturbations of the scalar field were suppressed. Both of these effects tend to suppress the density perturbations produced at the beginning of inflation, and they would be able to account for the suppression of the large-angle CMB anisotropy ob-

served by WMAP.

The $\delta\rho/\rho$ graphs present a small displacement with respect to N , when compared with the results obtained in Einstein's GR theory. This would change the constraint on the value of the parameters that appears in the scalar potentials $V(\sigma)$. In this way, we have shown that closed inflationary universe models in a JBD theory are less restricted than one analogous in Einstein's GR theory due to the introduction of a new parameter, i.e. the ω parameter. The inclusion of this parameter gives us a freedom that allows us to modify the cosmological model by simply modifying the corresponding value of this parameter.

As Linde mentioned in Ref.[4], the fine - tuning required for the construction of a closed inflationary model is much easier to achieve than the incredible fine - tuning required for the explanation of the enormously large mass and entropy of the universe, its homogeneity and isotropy, and the observed anisotropy of CMB without using inflation.

The models that we have treated are fine-tuned, such that the flatness of the universe remains a prediction of these inflationary scenarios. Nevertheless one should keep in mind that with the fine-tuning at a level of about one percent one can obtain a semi-realistic model of an inflationary universe with $\Omega > 1$ Ref.[4].

Acknowledgments

SdC was supported from COMISION NACIONAL DE CIENCIAS Y TECNOLOGIA through FONDECYT N^os 1030469, 1040624 and 1051086 grants. Also, it was partially supported by PUCV by grant 123.764.

-
- [1] H. V. Peiris et al. astro-ph/0302225; D. Spergel et al., astro-ph/0302209.
 - [2] C. L. Bennett et al., astro-ph/0302207.
 - [3] G. Efstathiou astro-ph/0303127.
 - [4] A. Linde, JCAP 002, 0305 (2003).
 - [5] M Sasaki, T. Tanaka, K. Yamamoto and J. Yokoyama, Phys. Lett. **317B**, 510 (1993).
 - [6] M. Bucher *et al*, Phys. Rev. D **52**, 3314 (1995); M. Bucher and N. Turok, Phys. Rev. D **52**, 5538 (1995).
 - [7] A. Linde, Phys. Lett. B, **351**, 99 (1995); A. Linde and A. Mezhlumian, Phys. Rev. D **52**, 6789 (1995).
 - [8] A. Linde, Phys. Rev. D **59**, 023503 (1999); A. Linde, M. Sasaki and T. Tanaka, Phys. Rev. D **59**, 123522 (1999).
 - [9] J.R. Gott, Nature **295**, 304 (1982).
 - [10] J.R. Gott and T.S. Statler, Phys. Lett. **136B**, 157 (1984).
 - [11] U. Moschella and R. Schaeffer, Phys. Rev. D **57**, 2147 (1998).
 - [12] S. del Campo and R. Herrera, Phys. Rev. D **67**, 063507 (2003).
 - [13] M. Bouchmadi and P. F. González-Días, Phys. Rev. D **65**, 063510 (2002); S. del Campo, R. Herrera and J. Saavedra, Phys. Rev. D **70**, 023507 (2004).
 - [14] M. White and D. Scott, Astrophys J. **459**, 415 (1996).
 - [15] G. Ellis, W. Stoerger, P. McEwan and P. Dunsby, Gen. Rel. Grav. **34**, 1445 (2002); G. Ellis, W. Stoerger, P. cEwan and P. Dunsby, Gen. Rel. Grav. **34**, 1461 (2002); S. del Campo, R. Herrera and J. Saavedra, Int. J. Mod. Phys. D **14**, 1, (2005).
 - [16] P. Jordan, Z. Phys. **157**, 112 (1959); C. Brans and R. Dicke, Phys. Rev. **124**, 925 (1961).
 - [17] A. Linde, Phys. Rev. **49**, 748 (1994).
 - [18] A. Linde, Phys. Lett. B **238**, 160 (1990).
 - [19] S. del Campo and A. Vilenkin, Phys. Rev. D **40**, 688 (1989); A. Linde, Phys. Lett B **249**, 18 (1990).
 - [20] T. Damour and K. Nordtvedt, Phys. Rev. Lett. **70**, 2217 (1993); Phys. Rev. D **48**, 3436 (1993).
 - [21] R. Nagata, T. Chiba and N. Sugiyama, Phys. Rev. D **69**, 083512 (2004).
 - [22] R. Nagata, T. Chiba and N. Sugiyama, Phys. Rev. D **66**, 103510 (2002); X. Chen and M. Kamionkowski, Phys. Rev. D **60**, 104036 (1999).
 - [23] A. Linde, Phys. Lett B **129**, 177 (1983).
 - [24] K. Freese, J. A. Frieman and V. Olinto, Phys. Rev. Lett. **65**, 3233 (1990).

- [25] F. C. Adams, J. Bond, K. Freese, J. A. Frieman and V. Olinto, Phys. Rev. D **47**, 426 (1993).
- [26] T. Banks, M. Dine, P. J. Fox and E. Gorbatov, JCAP **0306**, 001 (2003); A. Linde, hep-th/0503195.
- [27] A. Starobinsky and J. Yokoyama, gr-qc/9502002.